

Supplementary information

**BMS-708163 and Nilotinib restore synaptic dysfunction in human embryonic stem cell-derived
Alzheimer's disease models**

Hisae Nishioka^{1,2}, Norie Tooi¹, Takehisa Isobe¹, Norio Nakatsuji^{1,3}, Kazuhiro Aiba^{1, 4*}

¹Institute for Integrated Cell-Material Sciences (WPI-iCeMS), Kyoto University, Kyoto 606-8501,
Japan

²Graduate School of Medicine, Kyoto University, Kyoto 606-8501, Japan

³Institute for Frontier Medical Sciences, Kyoto University, Kyoto 606-8507, Japan

⁴Current address: Stem Cell & Device laboratory, Inc., Kyoto 600-8491, Japan

*Corresponding Author: Kazuhiro Aiba


Institute for Integrated Cell-Material Sciences (WPI-iCeMS), Kyoto University

Yoshida-Ushinomiya-cho, Sakyo-ku, Kyoto 606-8501, Japan

Tel: +81-75-753-9762


Fax: +81-75-753-9785

E-mail: kaiba@icems.kyoto-u.ac.jp




Supplementary Figure S1. **PS1-G378E neurons did not show significant differences in RAB3A and SV2B gene expression levels**

Gene expression levels of RAB3A, SV2B, Synaptophysin (SYP) and Synaptotagmin 1 (SYT1) were indicated in PS1-G378E neurons (G378E) relative to PS1-WT neurons (WT). β -actin was used as an internal control. Each gene expression level in PS1-WT neurons was defined as 1.0. Mann–Whitney U test was used to check for differences in expression levels. Four independent experiments, each time in triplicates were performed ($n = 4$). Mean \pm SD.



Supplementary Figure S2. **Cholinergic neurons derived from PS1-overexpressing hESCs.**


Immunocytochemistry using antibodies against a cholinergic neuron marker, choline acetyltransferase (ChAT, red) and a neuron marker, β III-tubulin (β III, green) were carried out. Cells were counterstained with 4',6-diamidino-2-phenylindole (DAPI, blue) to visualize nuclei. Scale bar, 100 μ m. A few cells ($0.9 \pm 0.7\%$) were detected as choline acetyltransferase-positive neurons in hESC-derived neurons used in this study.



Supplementary Figure S3. Drug responses and cell viability of PS1-G378E neurons derived from three subclones expressing mutant-PS1

The effects of A β inhibition (A) and cell survival (B) in the presence of various concentrations of DAPT using three subclones of PS1-G378E neurons. The amount of A β 40 and cell viability in DMSO-treated PS1-G378E neurons was defined as 1.0. Three independent experiments, each time in triplicates were performed (n = 3). Mean \pm SD.

Using 3 subclones (No.1 – 3) of the hESCs overexpressing PS1-G378E (PS1-G378E hESCs), we investigated whether there is any variation among neurons derived from the PS1-G378E hESC subclones. In our previous report, it was confirmed that the PS1 protein expression levels were not significantly different between different subclones¹. DAPT-response experiments showed that there were also no differences in DAPT responses (A β 40 reduction and cell viability) among the subclones (Supplementary Fig. S2). These data indicate that there are no significant variations among PS1-G378E subclones. In addition, all of clones had an identical genetic background because the site-specific gene integration method was applied for establishing clones overexpressing mutant PS1. Hence we used each single clone of PS1-G378E neurons for the experiments in this study.



Supplementary Figure S4. **Preparation of synaptosomes from the PS1-WT and PS1-G378E neurons**

Neural differentiation, following synaptosome preparation were independently carried out four times (n = 4). Immunoblot analyses of a pre-synaptic protein, SNAP25 were performed using the whole cells and synaptosomes of PS1-WT and PS1-G378E neurons. These data indicated that isolation of synaptosomes was successfully carried out in all preparation. β -actin (ACT) was used as an internal control. WT, PS1-wild type neurons; G378E, PS1-G378E neurons; wh, whole cells; sy, synaptosomes.

Supplementary Table S1. **Result of chemical screening**

| | Chemicals | Relative A β 40 ratios* | | Relative viability ratios (2 nd screening)** |
|--|--|-------------------------------|------------------------------|--|
| | | 1 st screening | 2 nd screening | |
| γ -secretase inhibitor | Semagacestat (LY-450139) ² | 0.088 | - | - |
| | Avagacestat (BMS-708163) ³ | 0.050 | - | - |
| | DAPT(GSI-IX) ⁴ | 0.14 | - | - |
| | LY-411575 ⁵ | 0.23 | - | - |
| | MK-0752 ⁶ | 0.10 | - | - |
| | YO-01027 (Dibenzazepine) ⁷ | 0.11 | - | - |
| Bcr-Abl inhibitor | Nilotinib ⁸ | 0.21 | 0.23 | 0.85 |
| Calcineurin inhibitor | Pimecrolimus | 0.20 | 0.27 | 0.87 |
| HMG-CoA reductase inhibitor | Fluvastatin Sodium ⁹ | 0.28 | 0.37 | 0.96 |
| | Rosuvastatin Calcium | 0.18 | 0.19 | 0.91 |
| Imidazole derivative | Sulconazole Nitrate salt | 0.23 | 0.25 | 1.04 |
| Selective estrogen receptor modulator | Toremifene Base | 0.27 | 0.23 | 0.92 |

*, The A β 40 level of DMSO treatment was considered to be 1.0.

**, The cell viability ratio of DMSO treatment was considered to be 1.0.

Supplementary Table S2. **The numerical data of figure 2d and 2e**

| | Relative A β 40 ratios (Mean \pm SD)* | | | | |
|----------------------|---|-----------------|-----------------|-----------------|-----------------|
| | K1 + chemicals | | | | |
| chemicals (μ M) | Nilo | Pime | Rosu | Sulc | Tore |
| 0.001 | 0.96 \pm 0.21 | 0.80 \pm 0.20 | 0.95 \pm 0.16 | 0.90 \pm 0.12 | 0.79 \pm 0.17 |
| 0.01 | 0.89 \pm 0.13 | 0.69 \pm 0.15 | 0.85 \pm 0.10 | 0.78 \pm 0.17 | 0.86 \pm 0.10 |
| 0.1 | 0.93 \pm 0.06 | 0.75 \pm 0.07 | 0.72 \pm 0.17 | 0.87 \pm 0.15 | 0.77 \pm 0.17 |
| 1 | 0.71 \pm 0.12 | 0.74 \pm 0.13 | 0.72 \pm 0.36 | 0.63 \pm 0.21 | 0.63 \pm 0.20 |
| 10 | 0.24 \pm 0.10 | 0.52 \pm 0.17 | 0.39 \pm 0.07 | 0.30 \pm 0.14 | 0.14 \pm 0.06 |

| | Relative viability ratios (Mean \pm SD)** | | | | |
|----------------------|---|-----------------|-----------------|-----------------|-----------------|
| | K1 + chemicals | | | | |
| chemicals (μ M) | Nilo | Pime | Rosu | Sulc | Tore |
| 0.001 | 1.16 \pm 0.10 | 1.05 \pm 0.09 | 0.97 \pm 0.18 | 1.04 \pm 0.19 | 1.19 \pm 0.25 |
| 0.01 | 1.19 \pm 0.23 | 1.07 \pm 0.15 | 0.93 \pm 0.11 | 1.19 \pm 0.22 | 1.02 \pm 0.09 |
| 0.1 | 1.09 \pm 0.09 | 0.99 \pm 0.11 | 0.94 \pm 0.12 | 0.98 \pm 0.19 | 1.05 \pm 0.19 |
| 1 | 1.05 \pm 0.18 | 1.02 \pm 0.10 | 0.84 \pm 0.18 | 1.11 \pm 0.12 | 0.97 \pm 0.22 |
| 10 | 0.62 \pm 0.14 | 0.86 \pm 0.22 | 0.72 \pm 0.14 | 0.72 \pm 0.12 | 0.67 \pm 0.22 |

*, The A β 40 level of DMSO treatment was considered to be 1.0.

**, The cell viability ratio of DMSO treatment was considered to be 1.0.

K1, KhES-1-derived neurons; Nilo, Nilotinib; Pime, Pimecrolimus; Rosu, Rosuvastatin Calcium;

Sulc, Sulconazole Nitrate; Tore, Toremifene Base.

References

- 1 Honda, M. *et al.* The modeling of Alzheimer's disease by the overexpression of mutant Presenilin 1 in human embryonic stem cells. *Biochem Biophys Res Commun* **469**, 587-592 (2016).
- 2 Lanz, T. A. *et al.* Concentration-Dependent Modulation of Amyloid- β in Vivo and in Vitro Using the γ -Secretase Inhibitor, LY-450139. *J Pharmacol Exp Ther* **319**, 924-933 (2006).
- 3 Gillman, K. W. *et al.* Discovery and Evaluation of BMS-708163, a Potent, Selective and Orally Bioavailable γ -Secretase Inhibitor. *ACS Med Chem Lett* **1**, 120-124 (2010).
- 4 Lanz, T. A. *et al.* The γ -Secretase Inhibitor N-[N-(3,5-Difluorophenacetyl)-l-alanyl]-S-phenylglycine t-butyl Ester Reduces A β Levels in Vivo in Plasma and Cerebrospinal Fluid in Young (Plaque-Free) and Aged (Plaque-Bearing) Tg2576 Mice. *J Pharmacol Exp Ther* **305**, 864-871 (2003).
- 5 Lanz, T. A. *et al.* Studies of A β Pharmacodynamics in the Brain, Cerebrospinal Fluid, and Plasma in Young (Plaque-Free) Tg2576 Mice Using the γ -Secretase Inhibitor N2-[(2S)-2-(3,5-Difluorophenyl)-2-hydroxyethanoyl]-N1-[(7S)-5-methyl-6-oxo-6,7-dihydro-5H-dibenzo[b,d]azepin-7-yl]-L-alaninamide (LY-411575). *J Pharmacol Exp Ther* **309**, 49-55 (2004).
- 6 Cook, J. J. *et al.* Acute γ -Secretase Inhibition of Nonhuman Primate CNS Shifts Amyloid Precursor Protein (APP) Metabolism from Amyloid- β Production to Alternative APP Fragments without Amyloid- β Rebound. *J Neurosci* **30**, 6743-6750 (2010).
- 7 Milano, J. *et al.* Modulation of Notch Processing by γ -Secretase Inhibitors Causes Intestinal Goblet Cell Metaplasia and Induction of Genes Known to Specify Gut Secretory Lineage Differentiation. *Toxicol Sci* **82**, 341-358 (2004).
- 8 Lonskaya, I. *et al.* Nilotinib-induced autophagic changes increase endogenous parkin level and ubiquitination, leading to amyloid clearance. *J Mol Med* **92**, 373-386 (2014).
- 9 Kurinami, H. *et al.* Prevention of amyloid beta-induced memory impairment by fluvastatin, associated with the decrease in amyloid beta accumulation and oxidative stress in amyloid beta injection mouse model. *Int J Mol Med* **21**, 531-537 (2008).